

The Value of a Multidisciplinary Integrated Approach on Improving the Quality of Care of Patients Affected by Obstructive Sleep Apnea Syndrome

Francesca Milano*

Private Practice of Orthodontics and Dental Sleep Medicine, Bologna, via Clavature 1, 40124 Bologna, Italy

Obstructive Sleep Apnea Syndrome (OSAS) is a common sleep respiratory disorder, characterized by snoring and partial (hypopnea) or complete (apnea) obstructions of airflow. Oxygen desaturation and sleep fragmentation, caused by the airflow limitations, can lead to an excessive *daytime sleepiness* and an increased risk of cardiovascular, cerebrovascular and metabolic diseases [1-8].

Nowadays, "Dental Sleep Medicine" plays an important role in the treatment of patients affected by OSAS. *Mandibular Advancing Devices* (MADs) represent a conservative, non-invasive and effective alternative for patients presenting a mild or moderate OSAS. Besides, it may give a valid possibility for patients affected by severe OSAS who refused CPAP (Continuous Positive Airway Pressure) and are not good candidates to surgery [9-11]. Furthermore, MADs can be combined to other treatment options, to obtain a better clinical outcome.

Involving a multidisciplinary expertise is essential for a correct *diagnosis and treatment plan*, and eventually the treatment management. Due to the high number of signs, symptoms and consequences of the syndrome, to the several etiological factors and the different treatment options, this becomes of paramount importance.

A comprehensive clinical examination should assess: -sleep-oriented present and past medical history;-daytime and nighttime symptoms;-findings of sleep testing; -comorbidities; -anthropometric characteristics; -findings of polysomnography;-anatomical and functional causes of obstruction; - morphology of skeletal and soft tissues; - oral (dental, periodontal and occlusal) and TMJ status. The objective is to create tailored *treatment plan*, which takes into account guidelines, contra/indications reported in literature, predictability of alternative therapies and preference of patients. This multidisciplinary approach gives the sleep dentist the following advantages: 1. improves patient *selection*; 2. reduces failures ratio, by means of *combined treatments*; 3. help to *control* the possible *persistent symptoms*.

Regarding *patients selection*, research on positive and negative predicting factors to MAD therapy is still a challenge because of the heterogeneity of causes, anatomical and functional characteristics and treatment outcomes. In attempt to identify good responders to MAD therapy, several authors emphasised the predictive role of the OSAS severity (good responders present a lower number of obstructive events at baseline polysomnography, when compared to poor responders), of the dependence on supine position (positional OSAS seems to present a better response to oral devices), of the BMI [12] (obesity is a negative predictive factor), of the gender (females normally obtain better results) and of the characteristics of the obstruction (a larger retrolingual airway caliber, a prevalent retropalatal obstruction and a lateral pattern of collapse seem to be positive predictive factors) [13]. Moreover, the evaluation of anatomical and functional dynamics of airway collapse during DISE (drug induced sleep endoscopy), can give valuable information [14,15].

Regarding *combined treatments*, lifestyle modifications, behavioral and positional therapy can influence MAD outcomes. Recently, Ravesloot et al. [16] have demonstrated the effect of a night positional trainer to control sleep posture. During the 23rd American Academy of

Dental Sleep Medicine Annual Meeting, Vanderveken et al. reported that by combining a night positional trainer to a MAD, achieved a significantly lower AHI (Apnea/Hypopnea Index), when compared to baseline or to only-MAD therapy results. However, when different anatomical sites are involved in the genesis of obstructive events, the combination of *MADs with ENT Surgery* can be needed [17]. Nevertheless, the treatment of high nasal resistance can improve MAD tolerability and outcome, by decreasing the tendency to mouth opening during sleep. Mouth opening often leads to an increased collapsibility of the pharynx and enhances the risk to lose the lower splint of the appliance (so the advancement of the mandible), especially when a low retention to teeth is present [18,19]. *Mandibular Advancing Devices* can be also *combined* to CPAP (Continuous Positive Airway Pressure), with the aim of reducing the effective airway pressure and the encumbrance of the interface in patients with poor compliance to CPAP [19].

Finally, the expertise of a neurologist can be essential to control *daytime sleepiness* that persists despite a significant improvement of the AHI, and *sleep quality and continuity*, thus improving the tolerance to the oral appliance.

On the other hand, the important role of an expert sleep dentist is: - to *recognize* symptoms of OSAS and to refer the patient to a sleep physician;- to *help in formulating* the complete *diagnosis* and the correct *treatment plan*, by evaluating the craniofacial morphology and the oral status (dental, periodontal and TMJ health); - to evaluate whether or not an oral device can be positioned;- to *carry out* the necessary periodontal, dental and prosthetic pretreatment;- to realize *MAD simulators* for sleep physician (used during polysomnography) and for ENT (used during DISE), for a higher predictability of the mandibular advancement effects [20-25];- to *complete* other treatments outcome, when it is considered insufficient [15]; - to collaborate with sleep physician in the evaluation of treatment outcome, taking into account the tolerance of the patient to mandibular advancement and his characteristics.

In conclusion, only a multidisciplinary approach to treat Obstructive Sleep Apnea Syndrome allows the best clinical outcome. "Coming together is a beginning, staying together is progress and working together is success. *Henry Ford*".

References

1. Bixler EO, Vgontzas AN, Lin HM, Ten Have T, Leiby BE, et al. (2000) Asso-

*Corresponding author: Francesca Milano, Private Practice of Orthodontics and Dental Sleep Medicine, Bologna, via Clavature 1, 40124 Bologna, Italy, Tel: +39 051 228084; Fax: +39 051 239889; E-mail: francescamilano@libero.it

Received November 18, 2013; Accepted December 07, 2013; Published December 09, 2013

Citation: Milano F (2013) The Value of a Multidisciplinary Integrated Approach on Improving the Quality of Care of Patients Affected by Obstructive Sleep Apnea Syndrome. J Interdiscipl Med Dent Sci 1: 104. doi: [10.4172/jimds.1000104](https://doi.org/10.4172/jimds.1000104)

Copyright: © 2013 Milano F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

- ciation of hypertension and sleep-disordered breathing. *Arch Intern Med* 160: 2289-2295.
2. Budhiraja R, Parthasarathy S, Quan SF (2007) Endothelial dysfunction in obstructive sleep apnea. *J Clin Sleep Med* 3: 409-415.
 3. Gottlieb DJ, Yenokyan G, Newman AB, O'Connor GT, Punjabi NM, et al. (2010) Prospective study of obstructive sleep apnea and incident coronary heart disease and heart failure: the sleep heart health study. *Circulation* 122: 352-360.
 4. Kario K (2009) Obstructive sleep apnea syndrome and hypertension: ambulatory blood pressure. *Hypertens Res* 32: 428-432.
 5. Neau JP, Paquereau J, Meurice JC, Chavagnat JJ, Gil R (2002) Stroke and sleep apnoea: cause or consequence? *Sleep Med Rev* 6: 457-469.
 6. Pamidi S, Tasali E (2012) Obstructive sleep apnea and type 2 diabetes: is there a link? *Front Neurol* 3: 126.
 7. Wang X, Bi Y, Zhang Q, Pan F (2013) Obstructive sleep apnoea and the risk of type 2 diabetes: a meta-analysis of prospective cohort studies. *Respirology* 18: 140-146.
 8. Guilleminault C, Stoohs R, Clerk A, Cetel M, Maistros P (1993) A cause of excessive daytime sleepiness. The upper airway resistance syndrome. *Chest* 104: 781-787.
 9. Kushida CA, Morgenthaler TI, Littner MR, Alessi CA, Bailey D, et al. (2006) Practice parameters for the treatment of snoring and Obstructive Sleep Apnea with oral appliances: an update for 2005. *Sleep* 29: 240-243.
 10. Schwarting S, Huebers U, Heise M, Schlieper J, Hauschild A (2007) Position paper on the use of mandibular advancement devices in adults with sleep-related breathing disorders. A position paper of the German Society of Dental Sleep Medicine (Deutsche Gesellschaft Zahnärztliche Schlafmedizin, DGZS). *Sleep Breath* 11: 125-126.
 11. Randerath WJ, Verbraecken J, Andreas S, Bettega G, Boudewyns A, et al. (2011) Non-CPAP therapies in obstructive sleep apnoea. *Eur Respir J* 37: 1000-1028.
 12. Marklund M, Stenlund H, Franklin KA (2004) Mandibular advancement devices in 630 men and women with obstructive sleep apnea and snoring: tolerability and predictors of treatment success. *Chest* 125: 1270-1278.
 13. Milano F, Mondini S, Billi MC, Gobbi R, Gracco A, et al. (2013) The impact of a multidisciplinary approach on response rate of mandibular advancing device therapy in patients with obstructive sleep apnoea syndrome. *Acta Otorhinolaryngol Ital* 33: 337-342.
 14. Johal A, Hector MP, Battagel JM, Kotecha BT (2007) Impact of sleep nasendoscopy on the outcome of mandibular advancement splint therapy in subjects with sleep-related breathing disorders. *J Laryngol Otol* 121: 668-675.
 15. Eichler C, Sommer JU, Stuck BA, Hörmann K, Maurer JT (2013) Does drug-induced sleep endoscopy change the treatment concept of patients with snoring and obstructive sleep apnea? *Sleep Breath* 17: 63-68.
 16. Ravesloot MJ, van Maanen JP, Dun L, de Vries N (2013) The undervalued potential of positional therapy in position-dependent snoring and obstructive sleep apnea-a review of the literature. *Sleep Breath* 17: 39-49.
 17. Millman RP, Rosenberg CL, Carlisle CC, Kramer NR, Kahn DM, et al. (1998) The efficacy of oral appliances in the treatment of persistent sleep apnea after uvulopalatopharyngoplasty. *Chest* 113: 992-996.
 18. the outcome of mandibular advancement splint therapy in subjects with sleep
 19. van Maanen JP, Richard W, Van Kesteren ER, Ravesloot MJ, Laman DM, et al. (2012) Evaluation of a new simple treatment for positional sleep apnoea patients. *J Sleep Res* 21: 322-329.
 20. Zeng B, Ng AT, Qian J, Petocz P, Darendeliler MA, et al. (2008) Influence of nasal resistance on oral appliance treatment outcome in obstructive sleep apnea. *Sleep* 31: 543-547.
 21. Vroegop AV, Vanderveken OM, Van de Heyning PH, Braem MJ (2012) Effects of vertical opening on pharyngeal dimensions in patients with obstructive sleep apnoea. *Sleep Med* 13: 314-316.
 22. El-Sohl AA, Moitheennazima B, Akinnusi ME, Churder PM, Lafornara AM (2011) Combined oral appliance and positive airway pressure therapy for obstructive sleep apnea: a pilot study. *Sleep Breath* 15: 203-208.
 23. Vanderveken OM, Vroegop AV, Van De Heyning PH, Braem MJ (2011) Drug-induced sleep endoscopy completed with a simulation bite approach for the prediction of the outcome of treatment of obstructive sleep apnea with mandibular repositioning appliances. *Oper Tech Otolaryngol* 22: 175-182.
 24. Almeida FR, Parker JA, Hodges JS, Lowe AA, Ferguson KA (2009) Effect of a titration polysomnogram on treatment success with a mandibular repositioning appliance. *J Clin Sleep Med* 5: 198-204.
 25. Gagnadoux F, Fleury B, Vielle B, Pételle B, Meslier N, et al. (2009) Titrated mandibular advancement versus positive airway pressure for sleep apnoea. *Eur Respir J* 34: 914-920.

Citation: Milano F (2013) The Value of a Multidisciplinary Integrated Approach on Improving the Quality of Care of Patients Affected by Obstructive Sleep Apnea Syndrome. *J Interdiscipl Med Dent Sci* 1: 104. doi: [10.4172/jimds.1000104](https://doi.org/10.4172/jimds.1000104)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/jimbt>